

CHEMICAL CONSTITUENTS AND
ANTIOXIDANT ACTIVITIES OF
AQUILARIA MALACCENSIS LAMK.
(AGARWOOD) LEAVES

NIK NOOR ASMA BINTI NIK WIL

MASTER OF SCIENCE
(INDUSTRIAL CHEMISTRY)

UNIVERSITI MALAYSIA PAHANG



SUPERVISOR'S DECLARATION

We hereby declare that We have checked this thesis and in our opinion, this thesis is adequate in terms of scope and quality for the award of the degree of Master of Science (Industrial Chemistry).

(Supervisor's Signature)

Full Name : DR. SAIFUL NIZAM BIN TAJUDDIN

Position : ASSOCIATE PROFFESOR

Date :

A handwritten signature in black ink, appearing to read 'Saiful'.

(Co-supervisor's Signature)

Full Name : NOR ADILA BINTI MHD OMAR

Position : LECTURER

Date :



STUDENT'S DECLARATION

I hereby declare that the work in this thesis is based on my original work except for quotations and citation which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Malaysia Pahang or any other institutions.

(Student's Signature)

Full Name : NIK NOOR ASMA BINTI NIK WIL

ID Number : MKD13003

Date :

CHEMICAL CONSTITUENTS AND ANTIOXIDANT ACTIVITIES OF
AQUILARIA MALACCENSIS LAMK. (AGARWOOD) LEAVES

NIK NOOR ASMA BINTI NIK WIL

Thesis submitted in fulfillment of the requirements
for the award of the degree of
Master of Science (Industrial Chemistry)

Faculty of Industrial Sciences & Technology
UNIVERSITI MALAYSIA PAHANG

AUGUST 2019

ACKNOWLEDGEMENT

Firstly, I would like to express my sincere gratefulness to Allah S.W.T for giving me a strength, inspiration, and wisdom to finish this thesis without any obstacles. All the experience and knowledge along my studies hopefully will give me the opportunity to develop my future career. I also hope this thesis can serve as references to other people who want guidance from this research study.

First and foremost, millions of thanks and highest appreciation dedicated to my supervisor, Assoc. Prof. Dr. Saiful Nizam Tajuddin for all his support, guidance and patience throughout completing this project. He also helps me in giving some advices and motivates in everything about this research until achieve a main objective. I would like to extend my deepest gratitude to my Co-Supervisor Mrs Adila Mhd Omar who gave me a lot of knowledge in terms of the bioassay study. Not forgotten to Assoc. Prof. Dr. Nadeem and Dr. Seema who teaches me in all technique related to the spectroscopic study. All these people have graciously provided me with their expertise and time.

Also, a very special thank you to my research group mates and members of the staff faculty FIST and Bio Aromatic who have gladly given their time to help me throughout my research. Last but not least, my dearest appreciation and heartfelt thanks to my beloved parents and family for their everlasting prayer and support the strengthen me to complete this project. Lastly, I am deeply indebted to my wonderful husband and daughter for giving me a spirit to achieving my dream as a graduate student's master. Without their aid and support this project would not have made it far.

ABSTRAK

Spesies *Aquilaria* mengandungi pelbagai metabolit sekunder termasuk pelbagai sebatian fenolik yang dikatakan bertindak sebagai antioksidan yang sangat berkesan. Kajian terhadap sifat fitokimia dan antioksidan spesies *Aquilaria malaccensis* dari keluarga Thymelaeaceae telah dilakukan. Pengekstrakan fitokimia dari daun dilakukan dengan menggunakan kaedah pengekstrakan konvensional; pengekstrakan refluks menggunakan air dan penyerapan dengan metanol yang menghasilkan ekstrak air (WE) dan ekstrak metanol (ME). Pengekstrakan, pemeringkatan dan pemisahan campuran menggunakan beberapa teknik kromatografi (kromatografi kolum dan kromatografi lapisan nipis) telah digunakan dalam proses pemisahan sebatian tulen. Untuk mengenal pasti sebatian yang mudah meruap dalam *A. malaccensis* daun, dua teknik digunakan, iaitu, pengekstrakan mikro fasa pepejal (SPME) dan desorpsi haba langsung (DTD), kemudian dianalisis dengan menggunakan teknik gas kromatografi iaitu, GC-FID dan GC-MS. Sementara itu, kromatografi cecair ultra tinggi dengan kuadropol spektrometri jisim masa penerbangan (UPLC-QToF / MS) digunakan untuk menentukan sebatian kimia yang tidak mudah meruap di dalam tumbuhan tersebut. Potensi antioksidan daripada ekstrak *A. malaccensis* dikawal oleh kadar perangkap radikal bebas (DPPH) dan pengurangan kapasiti antioksidan melalui tembaga (CUPRAC). Perbezaan signifikasi adalah berdasarkan nilai p di mana $p < 0.05$ dianggap ada perbezaan dan sebaliknya. Beberapa komponen yang biasa ditemui adalah asid *n*-heksadekanoik, eudesmol dan oxo-agarospirol. Ekstrak air dan metanol, yang mana mengandungi jumlah kandungan fenolik yang tinggi (191.005 ± 0.002 dan 177.927 ± 0.001 mg dari GAEs/g ekstrak) menunjukkan kuasa penurunan dan menghapuskan aktiviti radikal bebas yang tinggi. Pemisahan ekstrak heksana membawa kepada pengasingan friedelanol dan pemisahan ekstrak diklorometana membawa kepada pengasingan friedelin. Struktur sebatian tulen tersebut dikenalpasti melalui kaedah seperti 1D (^1H , ^{13}C , DEPTQ), 2D (COSY, HSQC, HMBC) NMR, MS, UV, FTIR dan juga melalui perbandingan maklumat spektra daripada kajian lepas.

ABSTRACT

Aquilaria species contains variety of secondary metabolites including various phenolic compounds which have been reported as excellent antioxidants. A study on phytochemical and antioxidant properties of *Aquilaria malaccensis* from Thymelaeaceae family was performed. The extraction of phytochemicals from the leaves were performed using conventional extraction methods: reflux extraction using water and maceration using methanol to obtain water extract (WE) and methanol extract (ME), respectively. Solvent–solvent extraction, fractionation and separation using different chromatographic techniques (column chromatography and thin layer chromatography) were used for isolation of pure compounds. In order to identify the volatile aroma compounds in the leaves of *A. malaccensis*, two methods were used namely solid-phase microextraction (SPME) and direct thermal desorption (DTD), in combination with gas chromatography (GC) and gas chromatography-mass spectrometry (GC-MS) analysis. The ultra-high-performance liquid chromatography-quadrupole time-of-flight mass spectrometry (UPLC-QToF/MS) was used for determination of non-volatile chemical compounds present in the leaves. The potential antioxidative activity of the leaves extracts of *A. malaccensis* was evaluated via radical scavenging assay (DPPH) and copper reducing antioxidant capacity (CUPRAC) assays. Significant differences were based on p values where $p < 0.05$ were considered significantly different and vice-versa. Some of the commonly identified chemical components were *n*-hexadecanoic acid, eudesmol and oxo-agarospirol. WE and ME extracts with the highest total phenolic contents (191.005 ± 0.002 and 177.927 ± 0.001 mg of GAEs/g extract) showed strong reducing power and scavenging radical activity. Friedelanol and friedelin were isolated from the fractionation of hexane and dichloromethane extracts, respectively. The structure of the isolated compounds was elucidated by using spectroscopic methods namely 1D (^1H , ^{13}C , DEPTQ), 2D (COSY, HSQC, HMBC) NMR, MS, UV, FTIR and by comparison with literature values of published data spectra.

TABLE OF CONTENT

DECLARATION	
TITLE PAGE	
ACKNOWLEDGEMENT	ii
ABSTRAK	iii
ABSTRACT	iv
TABLE OF CONTENT	v
LIST OF TABLES	ix
LIST OF FIGURES	x
LIST OF SYMBOLS	xi
LIST OF ABBREVIATION	xii
CHAPTER 1 INTRODUCTION	1
1.1 Introduction	1
1.2 Problem Statement	4
1.3 Research Objectives	4
1.4 Scope of Study	5
CHAPTER 2 LITERATURE REVIEW	6
2.1 Introduction	6
2.2 Botanical Overview	6
2.2.1 Thymelaeaceae	6
2.2.1 Genus <i>Aquilaria</i>	7
2.2.3 <i>Aquilaria malaccensis</i>	9
2.3 Medicinal Uses	11

2.4	Phytochemical Constituents of Genus <i>Aquilaria</i>	11
2.5	Method of Extraction	17
2.5.1	Solid-phase Microextraction (SPME)	17
2.5.2	Direct Thermal Desorption (DTD)	17
2.6	Methods of Analysis	18
2.7	Biological Activities of <i>Aquilaria</i> Leaves	19
2.7.1	Antioxidant Activity	20
CHAPTER 3 METHODOLOGY		24
3.1	Introduction	24
3.2	Materials	24
3.2.1	Plant sample	24
3.2.2	Materials for Extraction and Isolation	24
3.2.3	Plant Sample Preparation	25
3.3	Extraction Methods	25
3.3.1	Reflux Extraction	25
3.3.2	Maceration	25
3.3.3	Liquid-liquid Extraction	26
3.3.4	Solid-phase Microextraction (SPME)	28
3.3.5	Direct Thermal Desorption (DTD)	29
3.4	Phytochemical Analysis	30
3.4.1	Gas Chromatography-Mass Spectrometry (GC-MS)	30
3.4.2	Gas Chromatography-Flame Ionisation Detector (GC-FID)	30
3.4.3	Quadrupole Time-of-Flight Liquid Chromatography-Mass Spectrometry (QTOF-LCMS)	31
3.4.4	Fourier Transform Infrared Spectroscopy (FTIR)	31
3.4.5	UV-Vis Spectrophotometer (UV-Vis)	32

3.4.6	Nuclear Magnetic Resonance (NMR)	32
3.5	Total Phenolic Content (TPC)	32
3.6	Bioactivity Assay	33
3.6.1	DPPH radical-scavenging activity	33
3.6.2	Cupric Reducing Antioxidant Capacity (CUPRAC)	33
3.7	Separation and Purification	34
3.7.1	Compound I	34
3.7.2	Compound II	35
3.7	Statistical Analyses	35
CHAPTER 4 RESULTS & DISCUSSION		36
4.1	Introduction	36
4.2	Extraction	36
4.3	Profiling of <i>A. malaccensis</i> leaves extracts using GC and UPLC-QToF/MS	37
4.3.1	Profiling of Volatile Chemical Constituents by using SPME/DTD using GCMS/FID	37
4.3.2	Profiling of <i>A. malaccensis</i> Compounds using UPLC-QToF/MS	43
4.4	Total Phenolic Contents (TPC)	47
4.4.1	TPC of WE and its fraction	47
4.4.2	TPC of ME and its fraction	47
4.5	Bioactivities	49
4.5.1	DPPH and CUPRAC assay of WE and its fraction	49
4.5.2	DPPH and CUPRAC assay of ME and its fraction	51
4.6	Correlation of bioactivities with total phenol content	53
4.7	Compound isolation	54
4.7.1	Compound I (Friedelanol)	55
4.7.2	Compound II (Friedelin)	57

4.7.3 Bioactivity of isolated compound	60
CHAPTER 5 CONCLUSION	62
5.1 Conclusions	62
5.1 Recommendations	63
REFERENCES	64
APPENDIX A	73
APPENDIX B	74
APPENDIX C	75
APPENDIX D	76
APPENDIX E	77
APPENDIX F	79
APPENDIX G	88
APPENDIX H	91

LIST OF TABLES

Table 1.1	Pharmacological activities of agarwood plant	2
Table 2.1	Taxonomic classification of <i>Aquilaria malaccensis</i>	6
Table 2.2	Distribution of <i>Aquilaria</i> species in Malaysia	8
Table 2.3	Chemical constituents in different extracts of <i>Aquilaria</i> leaves	12
Table 2.4	Reported biological activities of crude leaves extracts from various species of <i>Aquilaria</i>	19
Table 3.1	Fractions from column chromatography of <i>A. malaccensis</i> hexane extract	34
Table 3.2	Fractions from column chromatography of <i>A. malaccensis</i> dichloromethane extract	35
Table 4.1	Tentative identification of volatile chemical compounds in WE and ME obtained from the leaves of <i>A. malaccensis</i> by using SPME	38
Table 4.2	Tentative identification of volatile chemical compounds in WE and ME obtained from the leaves of <i>A. malaccensis</i> by using DTD	40
Table 4.3	Percentage of peak area of chemical compounds in WE and ME obtained from the leaves of <i>A. malaccensis</i>	42
Table 4.4	Quantitative analysis of phenolic contents in WE and its fractions obtained from the leaves of <i>A. malaccensis</i>	47
Table 4.5	Quantitative analysis of phenolic contents in ME and its fractions obtained from the leaves of <i>A. malaccensis</i>	48
Table 4.6	IC ₅₀ in DPPH assay for WE and its fractions	51
Table 4.7	IC ₅₀ in DPPH assay for ME and its fractions	52
Table 4.8	Total phenolic content (TPC) and antioxidant activities (DPPH and CUPRAC) of <i>A. malaccensis</i>	54
Table 4.9	¹ H-NMR and ¹³ C-NMR data of friedelanol in CDCl ₃	56
Table 4.10	¹ H-NMR data of friedelin in CDCl ₃	58
Table 4.11	Results of DPPH assay for isolated compounds	61

LIST OF FIGURES

Figure 2.1	Part of <i>A. malaccensis</i> (1) Flower (2) Fruit	9
Figure 2.2	<i>A. malaccensis</i> (1) flowering branch, (2) flower, (3) longitudinal section of flower, (4) fruit and (5) longitudinal section of fruit	10
Figure 3.1	Extraction of <i>A. malaccensis</i> leaves and the extract fractionation by reflux extraction	26
Figure 3.2	Extraction of <i>A. malaccensis</i> leaves and the extract fractionation by maceration	27
Figure 3.3	Arrangement of apparatus in SPME sampling technique	28
Figure 3.4	Schematic diagram of SPME extraction method	29
Figure 3.5	Direct thermal desorption (DTD) adaptor	30
Figure 4.1	UPLC-QToF/MS positive BPI chromatogram of WE from the leaves of <i>A. malaccensis</i>	45
Figure 4.2	Major compounds plotted on (+) ESI-MS of WE	45
Figure 4.3	UPLC-QToF/MS positive BPI chromatogram of ME from the leaves of <i>A. malaccensis</i>	46
Figure 4.4	Major compounds plotted on (+) ESI-MS of ME	46
Figure 4.5	The DPPH free radical scavenging activity of <i>A. malaccensis</i> from WE and its fractions	50
Figure 4.6	Cupric ion reducing antioxidant capacity of WE and its fractions of <i>A. malaccensis</i>	51
Figure 4.7	The DPPH free radical scavenging activity of <i>A. malaccensis</i> from ME and its fractions	52
Figure 4.8	Cupric ion reducing antioxidant capacity of ME and its fractions of <i>A. malaccensis</i>	53
Figure 4.9	Structure of friedelanol	55
Figure 4.10	Structure of friedelin	57
Figure 4.11	EIMS spectrum data of friedelin	59
Figure 4.12	Mass fragmentation patterns of friedelin	60

LIST OF SYMBOLS

α	alpha
β	beta
$^{\circ}\text{C}$	Celsius
cm	centimetre
cm^{-1}	per centimetre
CuCl_2	copper (II) chloride
δ	chemical shift in ppm
eV	electron volt
γ	gamma
g	gram
hr	hour
KBr	potassium bromide
kg	kilogram
Kv	kilo volt
λ_{max}	maximum wavelength
μM	micromolar
μ	micro
μL	microlitre
m	metre
mL	millilitre
mm	millimetre
mM	millimolar
μM	micromolar
mg/mL	milligram per millilitre
MgCl_2	magnesium chloride
MHz	mega Hertz
m/z	mass to charge ratio
Na	sodium
NH_4OH	ammonium hydroxide
nm	nanometre
%	percentage

LIST OF ABBREVIATION

AA	ascorbic acid
BPI	based peak ion
^{13}C NMR	carbon nuclear magnetic resonance
CDCl_3	deuterated chloroform
COSY	correlation spectroscopy
CUPRAC	cupric reducing capacity
dbh	diameter at breast height
DCM	dichloromethane
DEPTQ	distortionless enhancement by polarization transfer with retention of quaternaries
DPPH	2,2-diphenyl-1-picrylhydrazyl
DTD	direct thermal desorption
EIMS	electron impact mass spectrometry
EtOAc	ethyl acetate
EtOH	ethanol
GC-FID	gas chromatography-flame ionization detector
GC-MS	gas chromatography-mass spectrometry
Hx	hexane
HPLC	high performance liquid chromatography
^1H NMR	proton nuclear magnetic resonance
HMBC	heteronuclear multiple bond coherence
HSQC	heteronuclear single quantum coherence
EC_{50}	concentration required to obtain a 50 % antioxidant effect
GAE	mg/g gallic acid equivalents
IC_{50}	concentration of drug required to inhibit cell growth by 50 %
IR	infrared
J	coupling constant
M^+	molecular ion
ME	methanol extract
MS	mass spectrum
NIST	National Institute of Standards and Technology

NMR	nuclear magnetic resonance
pH	power of hydrogen
RI	retention index
RP-18	reverse phase silica gel
SPME	solid phase microextraction
t_R	retention time
TIC	total ion current chromatogram
TPC	total phenolic content
TLC	thin layer chromatography
UPLC-QToF/MS	ultra-high-performance liquid chromatography-quadrupole time-of-flight mass spectrometry
UV-Vis	ultraviolet-visible spectroscopy
WE	water extract

REFERENCES

- Abkenar, D., ShivaYamini, Y., Shemirani, F., & Assadi, Y. (2006). Solid phase microextraction with gas chromatography-mass spectrometry: A very rapid method for identification of volatile organic compounds emitted by *Carum copticum*. *Natural Product Research*, 20(9), 850–859.
- Adam, A. Z., Tajuddin, S. N., Sudmoon, R., Chaveerach, A., Abdullah, U. H., Mahat, M. N., & Mohamed, R. (2018). Chemical constituents and toxicity effects of leaves from several agarwood tree species (*Aquilaria*). *Journal of Tropical Forest Science*, 30(3), 342–353.
- Akter, S., Islam, M. T., Zulkefeli, M., & Khan, S. I. (2013). Agarwood Production - A Multidisciplinary Field to be Explored in Bangladesh. *International Journal of Pharmaceutical and Life Sciences*, 2(1), 22–32.
- Alam, J., Mujahid, M., Badruddeen, Jahan, Y., Bagga, P., & Rahman, M. A. (2017). Hepatoprotective potential of ethanolic extract of *Aquilaria agallocha* leaves against paracetamol induced hepatotoxicity in SD rats. *Journal of Traditional and Complementary Medicine*, 7(1), 9–13.
- Ali, N. A. M., Jin, C. B., & Jamil, M. (2015). Agarwood (*Aquilaria malaccensis*) oils. In Victor R. Preedy (Eds)... *Essential Oils in Food Preservation, Flavor and Safety*. 173-180.
- Antonopoulou, M., Compton, J., Perry, L. S., & Al-Mubarak, R. (2010). The Trade and Use of Agarwood (Oudh) in the United Arab Emirates. *TRAFFIC Southeast Asia*.
- Apak, R., Guclu, K., Ozyurek, M., & Karademir, S. E. (2004). Novel Total Antioxidant Capacity Index for Dietary Polyphenols and Vitamins C and E, Using Their Cupric Ion Reducing Capability in the Presence of Neocuproine: CUPRAC Method. *Journal of Agricultural and Food Chemistry*, 52(26), 7970–7981.
- Aparna, V., Dileep, K. V., Mandal, P. K., Karthe, P., Sadasivan, C., & Haridas, M. (2012). Anti-Inflammatory Property of n-Hexadecanoic Acid: Structural Evidence and Kinetic Assessment. *Chemical Biology & Drug Design*, 80(3), 434–439.
- Aromdee, C., Pannangpetch, P., & Pranakhon, R. (2015). Effects of iriflophenone 3-C- β -glucoside on fasting blood glucose level and glucose uptake. *Pharmacognosy Magazine*, 11(41), 82-89.
- Azziz, S. S. S. A., Arriffin, N. M., Alimon, H., Sukari, M. A., & Naz, H. (2013). Chemical study of *Aquilaria crassna*. *Chemistry of Natural Compounds*, 49(3), 575–576.
- Bahrani, H., Mohamad, J., Paydar, M., & Rothan, H. (2014). Isolation and characterisation of acetylcholinesterase inhibitors from *Aquilaria subintegra* for the treatment of Alzheimer's disease (AD). *Curr Alzheimer Res*, 11(2), 206–214.
- Barden, A., Anak, N., Mulliken, T., & Song, M. (2000). Heart of the Matter: Agarwood use and trade and CITES implementation for *Aquilaria malaccensis*. Cambridge: TRAFFIC International, 1–52.
- Biesaga, M. (2011). Influence of extraction methods on stability of flavonoids. *Journal of Chromatography A*, 1218(18), 2505–2512.

- Blanchette, R. A., Heuveling, V. B (2009). Cultivated agarwood. *U.S. Patent No. 7,638,145. Washington, DC. U.S. Patent and Trademark Office.*
- Brand-Williams, W., Cuvelier, M. E., & Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT - Food Science and Technology*, 28(1), 25–30.
- Chakrabarty, K., Kumar, A., & Menon, V. (1994). Trade in Agarwood: A Study by TRAFFIC India.
- Chanda, S., & Dave, R. (2009). In vitro models for antioxidant activity evaluation and some medicinal plants possessing antioxidant properties : An overview. *African Journal of Microbiology Research*, 3(13), 981–996.
- Chua, L. S. L. (2018). Agarwood (*Aquilaria malaccensis*) in malaysia. *A Paper Presented during NDF Workshop Case Studies, Mexico.*
- CITES. (2003). Annex 2. *Review of Significant Trade Aquilaria Malaccensis*, (November), 45–76. Retrieved from <https://www.cites.org/sites/default/files/eng/com/pc/14/E-PC14-09-02-02-A2.pdf>
- Constant, M., & Collier, J. (1997). Headspace gas chromatography profiles of fruit-flavored malt beverages using solid-phase microextraction. *Journal of the American Society of Brewing Chemists*, 55(3), 112–118.
- Cui, J., Guo, S., & Xiao, P. (2011). Antitumor and antimicrobial activities of endophytic fungi from medicinal parts of *Aquilaria sinensis*. *Journal of Zhejiang University Science B*, 12(5), 385–392.
- Dahham, S. S., Ahamed, M., B, K., Saghir, S., M., Alsuede, F., S., Iqbal, A., M., & Majid, A., M., S., A. (1997). Bioactive essential oils from *Aquilaria crassna* for cancer prevention and treatment. *Rev. Ecuat. Cancerol*, 4(1), 14–19.
- Dahham, S. S., Majid, A., Saghir, S. M., & Al suede, F. S. (2014). Antibacterial , Antifungal and Antioxidant Activities of *Aquilaria crassna*. *International Journal of Advances in Science and Technology*, 196–200.
- Dahham, S. S., Tabana, Y. M., Iqbal, M. A., Ahamed, M. B. K., Ezzat, M. O., Majid, A. S. A., & Majid, A. M. S. A. (2015). The anticancer, antioxidant and antimicrobial properties of the sesquiterpene β -caryophyllene from the essential oil of *Aquilaria crassna*. *Molecules*, 20(7), 11808–11829.
- Dash, M., Patra, J. K., & Panda, P. P. (2008). Phytochemical and antimicrobial screening of extracts of *Aquilaria agallocha* Roxb ., 7(20), 3531–3534.
- Do, Q. D., Angkawijaya, A. E., Tran-Nguyen, P. L., Huynh, L. H., Soetaredjo, F. E., Ismadji, S., & Ju, Y. H. (2014). Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*, 22(3), 296–302.
- Donovan, D. G., & Puri, R. K. (2004). Learning from Traditional Knowledge of Non-timber Forest Products: Penan Benalui and the Autecology of *Aquilaria* in Indonesian Borneo. *Indonesian Borneo. Ecology and Society*, 9(3), 1–5.

- Duan, Z., Li, W., Dou, Z., Xie, H., He, A., & Shi, M. (2015). Extraction and antioxidant activity of flavonoids from *Aquilaria sinensis* (Lour.) Gilg leaves. *Food Science*, 36(6), 45–50.
- Escobedo-Martínez, C., Concepción Lozada, M., Hernández-Ortega, S., Villarreal, M. L., Gnecco, D., Enríquez, R. G., & Reynolds, W. (2012). ¹H and ¹³C NMR characterization of new cycloartane triterpenes from *Mangifera indica*. *Magnetic Resonance in Chemistry*, 50(1), 52–57.
- Falkovich, A. H., & Rudich, Y. (2001). Analysis of semivolatile organic compounds in atmospheric aerosols by direct sample introduction thermal desorption GC/MS. *Environmental Science and Technology*, 35(11), 2326–2333.
- Feng, J., & Yang, X. (2011). Liposolubility constituents from leaves of *Aquilaria sinensis*. *Journal of Chinese Materia Medica*, 36(15), 2092–2095.
- Feng, J., & Yang, X. (2012). Constituents from the leaves of *Aquilaria sinensis*. *Journal of Chinese Materia Medica*, 37(2), 230–234.
- Gao, Y., Liu, J., Lu, H., & Wei, Z. (2012). Two New 2- (2-Phenylethyl) chromen-4-ones from *Aquilaria sinensis* (Lour.) Gilg as sedative, analgesic, and digestive medicine in traditional Chinese medicine, 95, 951–954.
- Ghasemzadeh, A., & Ghasemzadeh, N. (2011). Flavonoids and phenolic acids: Role and biochemical activity in plants and human. *Journal of medicinal plants research*, 5(31), 6697-6703.
- Gunasekera, S. P., Kinghorn, A. D., Cordell, G. A., & Farnsworth, N. R. (1981). Plant anticancer agents. xix. constituents of *Aquilaria malaccensis*. *Journal of Natural Products*, 44(5), 569–572.
- Hara, H., Ise, Y., Morimoto, N., Shimazawa, M., Ichihashi, K., Ohyama, M., & Iinuma, M. (2008). Laxative Effect of Agarwood Leaves and Its Mechanism. *Bioscience, Biotechnology, and Biochemistry*, 72(2), 335–345.
- Hashim, Y. Z. H. Y., Kerr, G. P., Abbas, P., & Salleh, M. H. (2016). *Aquilaria* spp. (agarwood) as source of health beneficial compounds: A review of traditional use, phytochemistry and pharmacology. *Journal of Ethnopharmacology*, 189, 331–360.
- Hashim, Y.Z.H.Y., Abbas, P., & Awale, R. (2012). Antimicrobial Activities of Agarwood *A. subintegra* Leaves Extracts International Institute for Halal Research and Training (INHART), Ground Floor , Block, (July), 394–397.
- Hendra, H., Moeljopawiro, S., & Nuringtyas, T. R. (2016). Antioxidant and antibacterial activities of agarwood (*Aquilaria malaccensis* Lamk.) leaves. *AIP Conference Proceedings*, 1755.
- Huda, A., Munira, M., Fitrya, S., & Salmah, M. (2009). Antioxidant activity of *Aquilaria malaccensis* (thymelaeaceae) leaves. *Pharmacognosy Research*, 1(5), 270–273.
- Ishihara, M., Tsuneya, T., & Uneyama, K. (1991). Guaiane sesquiterpenes from agarwood. *Phytochemistry*, 30(10), 3343–3347.
- Ishihara, M., Tsuneya, T., & Uneyama, K. (1993). Components of the volatile concentrate of agarwood. *Journal of Essential Oil Research*, 5(3), 283–289.

- Ito, T., Kakino, M., Tazawa, S., Oyama, M., Maruyama, H., Araki, Y., ... Iinuma, M. (2012). Identification of Phenolic Compounds in *Aquilaria crassna* Leaves Via Liquid Chromatography-Electrospray Ionization Mass Spectroscopy. *Food Science and Technology Research*, 18(2), 259–262.
- Ito, T., Kakino, M., Tazawa, S., Watarai, T., Oyama, M., Maruyama, H., ... Iinuma, M. (2012). Quantification of polyphenols and pharmacological analysis of water and ethanol-based extracts of cultivated agarwood leaves. *J Nutr Sci Vitaminol (Tokyo)*, 58(2), 136–142.
- IUCN. (2019). *The IUCN Red List of Threatened Species 2018*. International Union for Conservation of Nature - IUCN. Retrieved from <https://www.iucnredlist.org/species/32056/2810130>. Accessed 7 Apr, 2019.
- Jiang, S., Yong Jiang, Guan, Y.-F., Tu, P.-F., Wang, K.-Y., & Chen, J.-M. (2011). Effects of 95% ethanol extract of *Aquilaria sinensis* leaves on hyperglycemia in diabetic db/db mice. *Journal of Chinese Pharmaceutical Sciences*, 20(6), 609–614.
- Kakino, M., Izuta, H., Ito, T., Tsuruma, K., Araki, Y., Shimazawa, M., ... Hara, H. (2010). Agarwood Induced Laxative Effects via Acetylcholine Receptors on Loperamide-Induced Constipation in Mice. *Bioscience, Biotechnology, and Biochemistry*, 74(8), 1550–1555.
- Kakino, M., Tazawa, S., Maruyama, H., Tsuruma, K., Araki, Y., Shimazawa, M., & Hara, H. (2010). Laxative effects of agarwood on low-fiber diet-induced constipation in rats. *BMC Complementary and Alternative Medicine*, 10 (1), 68.
- Kalia, K., Sharma, K., Singh, H. P., & Singh, B. (2008). Effects of Extraction Methods on Phenolic Contents and Antioxidant Activity in Aerial Parts of *Potentilla atosanguinea* Lodd. and Quantification of Its Phenolic Constituents by RP-HPLC. *Journal of Agricultural and Food Chemistry*, 56(21), 10129–10134.
- Kamonwannasit, S., Nantapong, N., Kumkrai, P., Luecha, P., Kupittayanant, S., & Chudapongse, N. (2013). Antibacterial activity of *Aquilaria crassna* leaf extract against *Staphylococcus epidermidis* by disruption of cell wall. *Annals of Clinical Microbiology and Antimicrobials*, 12(1), 1–7.
- Kang, Y. F., Chien, S. L., Wu, H. M., Li, W. J., Chen, C. T., Li, H. T., ... Chen, C. Y. (2014). Secondary metabolites from the leaves of *Aquilaria sinensis*. *Chemistry of Natural Compounds*, 50(6), 1110–1112.
- Khalil, A. S., Rahim, A. A. ., Taha, K. K. ., & Abdallah, K. B. (2013). Characterization of Methanolic Extracts of Agarwood Leaves. *Journal of Applied and Industrial Sciences*, 1(3), 78–88.
- Khan, N., & Mukhtar, H. (2008). Multitargeted therapy of cancer by green tea polyphenols. *Cancer Letters*, 269(2), 269–280.
- Khoddami, A., Wilkes, M., Roberts, T., Khoddami, A., Wilkes, M. A., & Roberts, T. H. (2013). Techniques for Analysis of Plant Phenolic Compounds. *Molecules*, 18(2), 2328–2375.
- Kim, S. J., Kim, J. A., Kim, D. H., Kwak, S. H., Yu, K. Y., Jang, S. Il, ... Jeong, S. Il. (2015). Bio-assay guided isolation and identification of α -glucosidase inhibitors

- from the leaves of *Diospyros lotus*. *Korean Journal of Pharmacognosy*, 46(2), 105–108.
- Konig, W. A., & Hochmuth, D. H. (2004). Enantioselective gas chromatography in flavor and fragrance analysis: Strategies for the identification of known and unknown plant volatiles. *Journal of Chromatographic Science*, 42(8), 423–439.
- Kovats, E. (1958). Gas chromatographic characterization of organic compounds. I. Retention indexes of aliphatic halides, alcohol, aldehyde and ketones. *Helv. Chim. Acta* 41: 1915–32.
- Kulip, J., Fan, L. N., Nurhuda Manshoor, Julius, A., Idris Said, Gisil, J., ... Tukip, W. F. (2010). Medicinal plants in Maliau Basin, Sabah, Malaysia. *Journal Of Tropical Biology And Conservation*, 6(January), 21–33.
- Kumar, S., & Pandey, A. K. (2013). Chemistry and Biological Activities of Flavonoids: An Overview. *The Scientif World J*, 4(2), 32–48.
- Lafrankie, J. V. (1994). Population Dynamics of Some Tropical Trees That Yield Non-Timber Forest Products. *Economic Botany*, 48(3), 301–309.
- Lee, N. Y., Yunus, M. A. C., Idham, Z., Ruslan, M. S. H., Aziz, A. H. A., & Irwansyah, N. (2016). Extraction and identification of bioactive compounds from agarwood leaves. *IOP Conference Series: Materials Science and Engineering*, 162(1).
- Lee, S. Y., & Mohamed, R. (2016). The Origin and Domestication of *Aquilaria*, an Important Agarwood-Producing Genus (pp. 1–20). Springer, Singapore.
- Lee, Y.-L., Huang, G.-W., Liang, Z.-C., & Mau, J.-L. (2007). Antioxidant properties of three extracts from *Pleurotus citrinopileatus*. *LWT - Food Science and Technology*, 40(5), 823–833.
- List, T. P. (2013). Thymelaeaceae. Retrieved January 12, 2019, from <http://www.theplantlist.org/1.1/browse/A/Thymelaeaceae/>. Accessed 7 Apr, 2019.
- Liu, Y., Chen, H., Yang, Y., Zhang, Z., Wei, J., Meng, H., ... Chen, H. (2013). Whole-tree agarwood-inducing technique: An efficient novel technique for producing high-quality agarwood in cultivated *Aquilaria sinensis* trees. *Molecules*, 18(3), 3086–3106.
- Liu, Y., Yang, X., & Liu, T. (2007). Yufeng, L., Xiuwei, Y., & Tonghua, L. (2007). GC-MS Analysis of Essential Oil from the Leaves of *Aquilaria sinensis* [J]. *Modern Chinese Medicine*, 8, 004.
- López-Sampson, A., & Page, T. (2018). History of Use and Trade of Agarwood. *Economic Botany*, 72(1), 107–129.
- Maisuthisakul, P., Suttajit, M., & Pongsawatmanit, R. (2007). Assessment of phenolic content and free radical-scavenging capacity of some Thai indigenous plants. *Food Chemistry*, 100(4), 1409–1418.
- Manuhara, G. J., Mentari, G. P., Khasanah, L. U., & Utami, R. (2018). Aqueous Extract Composition of Spent Ginger (*Zingiber officinale* var. *Amarum*) from Essential Oil Distillation. In *IOP Conference Series: Materials Science and Engineering*, 333 (1), IOP Publishing.

- Miniyar, P. B., Chitre, T. S., Karve, S. S., Deuskar, H. J., & Jain, K. S. (2008). Anti - oxidant activity of ethyl acetate extract of *Aquilaria agallocha* on nitrite - induced methemoglobin formation. *Int J Green Pharm*, 2(2), 116–117.
- Mohamed, R., Jong, P. L., & Kamziah, A. K. (2014). Fungal inoculation induces agarwood in young *Aquilaria malaccensis* trees in the nursery. *Journal of Forestry Research*, 25(1), 201–204.
- Monkodkaew, S., Loetchutinat, C., Nuntasaeen, N., & Pompimon, W. (2009). Identification and antiproliferative activity evaluation of a series of triterpenoids isolated from *Flueggea virosa* (Roxb. ex Willd.). *American Journal of Applied Sciences*, 6(10), 1800-1806.
- Molyneux, P. (2004). The use of the stable free radical diphenylpicryl-hydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin Journal of Science and Technology*, 26(2), 211–219.
- Moosa, S. (2010). Phytochemical and anti oxidant screening of *Aquilaria malaccensis* leaves. In *Proceedings of the Nuclear Malaysia R&D Seminar Proceedings of the Nuclear Malaysia R&D Seminar 12- Oct 2010. Bangi (Malaysia)*.
- Naef, R. (2011). The volatile and semi-volatile constituents of agarwood, the infected heartwood of *Aquilaria* species: A review. *Flavour and Fragrance Journal*, 26(2), 73–87.
- Ng, L. T., Chang, Y. S., & Kadir, A. A. (1997). A review on agar (gaharu) producing *Aquilaria* species. *Journal of Tropical Forest Products*, 2(2), 272–285.
- Nie, C., Song, Y., Chen, D., Xue, P., Tu, P., Wang, K., & Chen, J. (2009). Studies on chemical constituents of leaves of *Aquilaria sinensis*. *China Journal of Chinese Materia Medica*, 34(7), 858–860.
- Oktay, M. uni., Gulein, I., & Kufrevioglu, I. O. (2003). Determination of in vitro antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *LWT - Food Science and Technology*, 36(2), 263–271.
- Okudera, Y., & Ito, M. (2009). Production of agarwood fragrant constituents in *Aquilaria calli* and cell suspension cultures. *Plant Biotechnology*, 26(3), 307–315.
- Oyen, L. P. A. . N. X. D. (1999). *Plant Resources of South-East Asia No. 19 : Essential-oil plants*. the Netherlands: Leiden : Backhuys Publishers.
- Pandey, K. B., & Rizvi, S. I. (2009). Plant Polyphenols as Dietary Antioxidants in Human Health and Disease. *Oxidative Medicine and Cellular Longevity*, 2(5), 270–278.
- Pawliszyn, J. (2007). *Applications of solid phase microextraction*. Royal Society of Chemistry. Cambridge, UK. The Royal Society of Chemistry.
- PerezCoello, M. S., Sanz, J., & Cabezudo, M. D. (1997). Analysis of volatile components of oak wood by solvent extraction and direct thermal desorption gas chromatography mass spectrometry. *Journal of Chromatography A*, 778(1–2), 427–434.
- Persoon, G. A., & van Beek, H. H. (2008). Growing ‘The Wood of The Gods’: Agarwood Production in Southeast Asia (pp. 245–262). Springer, Dordrecht.

- Pranakhon, R., Pannangpetch, P., & Aromdee, C. (2011). Antihyperglycemic activity of agarwood leaf extracts in STZ-induced diabetic rats and glucose uptake enhancement activity in rat adipocytes. *Songklanakarin Journal of Science and Technology*, 33(4), 405–410.
- Prieto, P., Pineda, M., & Aguilar, M. (1999). Spectrophotometric Quantitation of Antioxidant Capacity through the Formation of a Phosphomolybdenum Complex: Specific Application to the Determination of Vitamin E. *Analytical Biochemistry*, 269(2), 337–341.
- Prior, R. L., Wu, X., & Schaich, K. (2005). Standarized Methods for the Determination of Antioxidant Capacity and Phenolics in Foods and Dietary Supplements. *Journal of Agricultural and Food Chemistry*, 53(10), 4290–4302.
- Prosen, H., & Zupancic-Kralj, L. (1999). Solid-phase microextraction. *TrAC Trends in Analytical Chemistry*, 18(4), 272–282.
- Qi, J., Lu, J., Liu, J., & Yu, B.-Y. (2009). Flavonoid and a rare benzophenone glycoside from the leaves of *Aquilaria sinensis*. *Chemical & Pharmaceutical Bulletin*, 57(2), 134–137.
- Rahman, H., Vakati, K., & Eswaraiah, M. C. (2012). In-vivo and in-vitro anti-inflammatory activity of *Aquilaria agallocha* oil. *International Journal of Basic Medical Sciences and Pharmacy*, 2(1), 7–10.
- Rahuman, A. A., Gopalakrishnan, G., Ghouse, B. S., Arumugam, S., & Himalayan, B. (2000). Effect of *Feronia limonia* on mosquito larvae. *Fitoterapia*, 71(5), 553–555.
- Ray, G., Leelamanit, W., Sithisarn, P., & Jiratchariyakul, W. (2014). Antioxidative Compounds from *Aquilaria crassna* Leaf. *Mahidol University Journal of Pharmaceutical Sciences*, 41(4), 54–58.
- Reiter, R. J. (1995). Oxidative processes and antioxidative defense mechanisms in the aging brain. *The FASEB Journal: Official Publication of the Federation of American Societies for Experimental Biology*, 9(7), 526–533.
- Richter, J., & Schellenberg, I. (2007). Comparison of different extraction methods for the determination of essential oils and related compounds from aromatic plants and optimization of solid-phase microextraction/gas chromatography. *Analytical and Bioanalytical Chemistry*, 387(6), 2207–2217.
- Roby, M. H. H., Sarhan, M. A., Selim, K. A.-H., & Khalel, K. I. (2013). Evaluation of antioxidant activity, total phenols and phenolic compounds in thyme (*Thymus vulgaris* L.), sage (*Salvia officinalis* L.), and marjoram (*Origanum majorana* L.) extracts. *Industrial Crops and Products*, 43, 827–831.
- Rodrigues, F., Caldeira, M., & Câmara, J. S. (2008). Development of a dynamic headspace solid-phase microextraction procedure coupled to GC–qMSD for evaluation the chemical profile in alcoholic beverages. *Analytica Chimica Acta*, 609(1), 82–104.
- Samadi, M., Abidin, Z. Z., Yunus, R., Awang Biak, D. R., Yoshida, H., & Lok, E. H. (2017). Assessing the kinetic model of hydro-distillation and chemical composition of *Aquilaria malaccensis* leaves essential oil. *Chinese Journal of Chemical*

- Engineering*, 25(2), 216–222.
- Sattayasai, J., Bantadkit, J., Aromdee, C., Lattmann, E., & Airara, W. (2012). Antipyretic, analgesic and anti-oxidative activities of *Aquilaria crassna* leaves extract in rodents. *Journal of Ayurveda and Integrative Medicine*, 3(4), 175-179.
- Sharma, O. P., & Bhat, T. K. (2009). DPPH antioxidant assay revisited. *Food Chemistry*, 113(4), 1202–1205.
- Shimada, K., Fujikawa, K., Yahara, K., & Nakamura, T. (1992). Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin emulsion. *Journal of Agricultural and Food Chemistry*, 40(6), 945–948.
- Shirey, R. E. (2000). Optimization of extraction conditions and fiber selection for semivolatile analytes using solid-phase microextraction. *Journal of chromatographic science*, 38(7), 279-288.
- Singleton, V. L., Orthofer, R., & Lamuela-Raventós, R. M. (1998). Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods in Enzymology*, 299(1974), 152–178.
- Singl Sushil Chandra Sati, Nitin Sati, U. R. and O. P. S. (2010). Medicinal Plants as a Source of Antioxidants. *Research Journal of Phytochemistry*, 4(4), 213–224.
- Tajuddin, S. N., & Yusoff, M. M. (2010). Chemical composition of volatile oils of *Aquilaria malaccensis* (Thymelaeaceae) from Malaysia. *Natural Product Communications*, 5(12), 1965–1968.
- Taskin, H., Baktemur, G., Kafkas, E., & Buyukalaca, S. (2013). Determinaton of Volatile Aroma Components of Gyromitra Mushroom Using Headspace Gas Dietary intake, Chromatography Mass Spectrometry (HS-GC/MS) Target hazard quotients, Ready-to-eat-foods,. *Turkish Journal of Agriculture - Food Science and Technology*, 1(1), 8–11.
- Tay, P. Y., Tan, C. P., Abas, F., Yim, H. S., & Ho, C. W. (2014). Assessment of extraction parameters on antioxidant capacity, polyphenol content, epigallocatechin gallate (EGCG), epicatechin gallate (ECG) and iriflophenone 3-C-β-glucoside of agarwood (*Aquilaria crassna*) young leaves. *Molecules*, 19(8), 12304–12319.
- Turkmen, N., Sari, F., & Velioglu, Y. S. (2006). Effects of extraction solvents on concentration and antioxidant activity of black and black mate tea polyphenols determined by ferrous tartrate and Folin–Ciocalteu methods. *Food Chemistry*, 99(4), 835–841.
- Ueda, J. ya, Imamura, L., Tezuka, Y., Tran, Q. L., Tsuda, M., & Kadota, S. (2006). New sesquiterpene from Vietnamese agarwood and its induction effect on brain-derived neurotrophic factor mRNA expression in vitro. *Bioorganic and Medicinal Chemistry*, 14(10), 3571–3574.
- Vernon H. Heywood, Richard K. Brummitt, A. C. and O. S. (2007). Flowering Plant Families of the World. Royal Botanic Gardens, Kew.
- Wardencki, W., Michulec, M., & Curyło, J. (2004). A review of theoretical and practical aspects of solid-phase microextraction in food analysis. *International Journal of Food Science and Technology*, 39(7), 703–717.

- Wetwitayaklung, P., Thavanapong, N., & Charoenteeraboon, J. (2009). Chemical Constituents and Antimicrobial Activity of Essential Oil and Extracts of Heartwood of *Aquilaria crassna* Obtained from Water Distillation and Supercritical Fluid Carbon Dioxide Extraction. *Silpakorn U Science & Technology Journal*, 3(1), 25–33.
- Wil, N. N. A., Mohd Omar, N. A., Awang Ibrahim, N., & Tajuddin, S. N. (2014). In vitro antioxidant activity and phytochemical screening of *Aquilaria malaccensis* leaf extracts. *Journal of Chemical and Pharmaceutical Research*, 6(12), 688–693.
- Yang, X. B., Feng, J., Yang, X. W., Zhao, B., & Liu, J. X. (2012). Aquisiflavoside, a new nitric oxide production inhibitor from the leaves of *Aquilaria sinensis*. *Journal of Asian Natural Products Research*, 14(9), 867–872.
- Yu, Q., Qi, J., Yu, H. X., Chen, L. L., Kou, J. P., Liu, S. J., & Yu, B. Y. (2013). Qualitative and quantitative analysis of phenolic compounds in the leaves of *Aquilaria sinensis* using liquid chromatography-mass spectrometry. *Phytochemical Analysis*, 24(4), 349–356.
- Zhang, L., Brockelman, W. Y., & Allen, M. A. (2008). Matrix analysis to evaluate sustainability: The tropical tree *Aquilaria crassna*, a heavily poached source of agarwood. *Biological Conservation*, 141(6), 1676–1686.
- Zhang, W., & Ni, B. (2011). GC-MS analysis of the chemical components of the volatile oil from leaves of *Aquilaria sinensis* (Lour.) Gilg. *Medicinal Plant*, 2(8), 34–36.
- Zhou, M., Wang, H., Suolangjiba, Kou, J., & Yu, B. (2008). Antinociceptive and anti-inflammatory activities of *Aquilaria sinensis* (Lour.) Gilg. Leaves extract. *Journal of Ethnopharmacology*, 117(2), 345–350.
- Zich, F., & Compton, J. (2002). The Final Frontier: Towards Sustainable Management of Papua New Guinea's Agarwood Resource. A *TRAFFIC Oceania Report in Conjunction with WWF South Pacific Programme*.
- Zulkifle, N. L., Omar, N. A. M., Tajudin, S. N., & Shaari, M. R. (2013). Antidiabetic Activities of Malaysian Agarwood (*Aquilaria spp.*) Leaves Extract. *Conference on Industry-Academia Joint Initiatives in Biotechnology CIA*.